1. Once more the RIKZ data: In the same data where ShannonW is given you also find a column "Richness" which is just the number of species found at each sample site. How does Richness depend on week, angle1, angle2, exposure, salinity, temperature, NAP, penetrability, grain-size, humus, chalk, sorting1 and Beach? Fit Poisson and quasipoisson models.

2. Collet<sup>1</sup> and Venables and Ripley<sup>2</sup> report an experiment to investigate the toxicity of transcyphemethrin to the tobacco budworm *Heliothis virescens*. Batches of 20 male and 20 female moths were exposed to different doses of trans-cyphemethrin for three days. The following table shows the numbers of dead or knocked down moths.

dose $[\mu g]$	1	2	4	8	16	32
male	1	4	9	13	18	20
female	0	2	6	10	12	16

How does the probability of a moth to be killed or knocked out depend on the dose and on the moth's sex? Fit a logistic regression model. Check whether the model fit can be improved by rescaling the dose in an appropriate way. Present your final results graphically with 95% confidence bands.

**3.** The data file TbDeerAndBoar.txt contains data from a survey of Vicente et al.<sup>3</sup>, see also Zuur et al.<sup>4</sup>. Boars and deers on 32 farms in Spain were tested for tubercolosis (Tb) and for the parasite *Elaphostrongylus cervi*. The table contains the numbers of sampled individuals and the numbers of positively tested individuals on each farm. The other variables in the table describe the habitat: the percentage of open land, pine and scrubs plantation, density of quercus plants, density of quercus trees, a wild boar abundance index, a red deer abundance index, the size of the habitat in ha, and whether the habitat was fenced (1) or not (0).

How does the risk of a red deer / wild boar to be infected with tubercolosis / *Elaphostrongylus cervi* depend on the other variables?

<sup>2</sup>Venables, W.N., Ripley, B.D (2002) *Modern Applied Statistics with S, 4th ed.* Springer, New York.

<sup>&</sup>lt;sup>1</sup>Collet, D. (1991) *Modelling Binary Data*. Chapman & Hall, London.

<sup>&</sup>lt;sup>3</sup>Vicente, J., Höfle, U., Garrido, J.M., Fernandez-de-Mera, I.G., Juste, R., Barralb, M., Gortarzar, C. (2006) Wild boar and red deer display high prevalences of tubercolosis-like lesions in Spain. *Veterinary Research* **37**: 107–119

<sup>&</sup>lt;sup>4</sup>Zuur, A.F., Ieno, E.N., Walker, N.J., Saveliev, A.A., Smith, G.M. (2009) *Mixed Effects Models and Extensions in Ecology with R.* Springer, New York.